

-1-

Date: October 23, 2001 Express Mail Label No. EV005366900 US

Inventors: Krzysztof B. Wicher, Olof Peder Holst, Maher Youssef Abou Hachem,  
Eva Margareta Nordberg Karlsson and Gudmundur O. Hreggvidsson

Attorney's Docket No.: 2739.2001-001

## THERMOSTABLE CELLULASE

### RELATED APPLICATIONS

This application is a continuation-in-part of and claims priority to PCT Application No. PCT/IS01/00012, filed 15 June 2001 (in English), which is a continuation-in-part of U.S. Application No. 09/594,884, filed June 15, 2000; <sup>now abandoned</sup> the entire teachings of the above applications are incorporated herein by reference.

### BACKGROUND OF THE INVENTION

Cellulases are enzymes that are capable of hydrolyzing cellulose. The products of the reaction include cellobiose and glucose which can in used for a variety of applications. For example, glucose obtained by cellulase catalyzed hydrolysis of plant cellulose can be fermented to produce ethanol which can be used as a fuel.

Cellulases can also be used in the de-inking and refining of recycled paper. Enzymes with high thermal stability are particularly useful in these applications because elevated temperatures are commonly used in these processes. Thermostable cellulases are also useful in the consumer products and food industries, for example, in extracting and clarifying juice from fruits or vegetables. Cellulases and particularly thermostable cellulases also have applications in the textile and laundry industries. For example, the enzyme can be used to remove microfibers from the surface of cotton garments (or other garments made of cellulose based fabric), thereby brightening the color and removing

the dull look that comes with wear. Cellulases are also useful for cleaning garments, for example as additives to detergents and for producing a "stone-washed" effect on indigo dyed denim (see, U.S. Patent No. 4,912,056).

The development of thermostable cellulases with improved stability and/or catalytic properties would provide advantages for the above-referenced applications and certain other applications. Therefore, a need exists for improved thermostable cellulases which can be easily produced.

#### SUMMARY OF THE INVENTION

The invention relates to polypeptides having thermostable cellulase activity.

10 The polypeptides of the invention are variants of full-length or naturally occurring proteins that have thermostable cellulase activity and are readily produced in large quantities by expression in a host cell such as *Escherichia coli*. In one embodiment, the polypeptide is a variant of a glycosyl hydrolase of family 12 wherein one or more of the amino acid residues that are not part of the catalytic domain (e.g, one or more amino

15 acid residues in the amino terminal hydrophobic domain and/or linker moiety) are deleted. In preferred embodiments, the polypeptide is derived from a thermophilic organism from a *Rhodotermus* species such as *R. marinus*, *R. obamae*, and *R. obamensis*; or a *Pyrococcus* species, including *P. abyssi*, *P. endeavori*, *P. furiosus*, *P. horikoshi*, *P. shinkai*, and *P. woesei*. In a more particular embodiment, the polypeptide

20 comprises the amino acid sequence of SEQ ID NO:2 wherein one or more of the amino acid residues from position one to about position 40 are deleted. In even more particular embodiments, the polypeptide can have an amino acid sequence selected from residues 18-261 of SEQ ID NO:2, residues 19-261 of SEQ ID NO:2, residues 20-261 of SEQ ID NO:2, residues 21-261 of SEQ ID NO:2, residues 22-261 of SEQ ID NO:2,

25 residues 23-261 of SEQ ID NO:2, residues 24-261 of SEQ ID NO:2, residues 25-261 of SEQ ID NO:2, residues 26-261 of SEQ ID NO:2, residues 27-261 of SEQ ID NO:2, residues 28-261 of SEQ ID NO:2, residues 29-261 of SEQ ID NO:2, residues 30-261 of SEQ ID NO:2, residues 31-261 of SEQ ID NO:2, residues 32-261 of SEQ ID NO:2,

residues 33-261 of SEQ ID NO:2, residues 34-261 of SEQ ID NO:2, residues 35-261 of SEQ ID NO:2, residues 36-261 of SEQ ID NO:2, residues 37-261 of SEQ ID NO:2 or residues 38-261 of SEQ ID NO:2.

The polypeptides of the invention can have improved catalytic properties and/or stability relative to full-length enzyme. In one embodiment, the polypeptide has a half-life of at least about 3.5 hours at 90°C. In another embodiment, the polypeptide has a specific activity that is at least about two times greater than the specific activity of a protein consisting of the amino acid sequence of SEQ ID NO:2.

In particularly preferred embodiments, the polypeptide has the amino acid sequence of residues 18-261 of SEQ ID NO:2 or the amino acid sequence of residues 38-261 of SEQ ID NO:2.

The invention also relates to isolated nucleic acids which encode a polypeptide of the invention and to constructs which comprises an isolated nucleic acid of the invention that is operatively linked to one or more regulatory sequences.

The invention also relates to host cells which comprise an isolated nucleic acid or construct of the invention, and to a method of producing a polypeptide having thermostable cellulase activity. In one embodiment, the method comprises maintaining a host cell of the invention under conditions suitable for expression of the polypeptide that has thermostable cellulase activity.

## BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing and other objects, features and advantages of the invention will be apparent from the following more particular description of preferred embodiments of the invention.

Fig. 1 is a graph illustrating the growth of *Escherichia coli* (*E. coli*) strain BL21(DE3) transfected with constructs encoding full-length Cel12A or  $\delta$ CEL12AH. Expression of Cel12A or  $\delta$ CEL12AH was induced by adding isopropyl  $\beta$ -D-thio-galactopyranoside (IPTG) to the cultures (time of addition of IPTG marked with the arrow). Cultures containing cells transfected with the construct encoding

full-length Cel12A were induced ( ) or not induced (○), and cultures containing cells transfected with the construct encoding  $\delta$ CEL12AH were induced (▲) or not induced (Δ). Induction of expression of full-length Cel12A (●) resulted in a decrease in the  $OD_{600}$ .

- 5 Fig. 2 is a photograph of an SDS-polyacrylamide gel stained with Coomassie brilliant blue G250. *E. coli* expression constructs encoding full-length Cel12A or variants thereof were induced and harvested 2.5 hours later. Crude cell contents were separated by SDS-PAGE and visualized by staining. Lane 1, uninduced culture; Lane 2, cells induced to express full-length Cel12A; Lane 3, cells induced to express  
10 Cel12A(sp-)H; Lane 4, cells induced to express  $\delta$ Cel12AH.

Figs. 3A-3D are schematic representations of full-length Cel12A (A) and variants  $\delta$ Cel12AH (B), Cel12A(sp-)H (C) and Cel12A(sp-) (D). S.P.- signal peptide. [EP]<sub>4</sub>-linker moiety. C.D.- catalytic domain. H.t.-HSV-Tag. H6- hexa-histidine.

- Fig. 4 is a graph showing the thermal unfolding curves for Cel12A(sp-)H (■)  
15 and  $\delta$ Cel12AH (Δ) measured by differential scanning calorimetry.

## DETAILED DESCRIPTION OF THE INVENTION

A description of preferred embodiments of the invention follows.

- Rhodothermus marinus* (*R. marinus*) is a thermophilic heterotrophic slightly halophilic marine eubacterium growing optimally at 65°C. *R. marinus* produces several  
20 thermostable glycosyl-hydrolases including a cellulase (Cel12A) (Hreggvidsson, G.O., *et al.*, *Applied and Environmental Microbiology*, 62:3047-3049 (1996)), which has been cloned in *Escherchia coli* (*E.coli*) and characterised (Halldorsdottir, S. *et al.*, *Appl Microbiol Biotechnol*, 49:277-284 (1998)). Cel12A, which belongs to family 12 glycosyl hydrolases, is among the most thermostable endo-cellulases, with maximal  
25 catalytic activity at temperatures substantially higher than the optimal temperature for growth of *R. marinus*. Many of cellulases in this family (family 12 glycosyl hydrolases) contain a highly hydrophobic signal peptide (Garda, A.L. *et al.*, *Biochem J.*,

324:403-411 (1997); Ooi, T. *et al.*, *Nucleic Acids Res.*, 18:5885 (1990)). Family 12 glycosyl-hydrolases catalyse the hydrolysis of  $\beta$ -1,4 glycosidic bonds primarily in mixed linkage (1-3),(1-4)  $\beta$ -D-glucans.

The invention relates to improved thermostable cellulases, nucleic acids and  
5 constructs (e.g., expression vectors) encoding the improved thermostable cellulases and to methods for producing the improved enzymes. As described herein, low yields of the thermostable cellulase Cel12A from *R. marinus* were obtained when a nucleic acid encoding the enzyme was expressed in *E. coli*. Low yields were obtained because the recombinant polypeptide was cytotoxic to the bacterial host cells. In addition, severe  
10 aggregation occurred when moderately high concentrations of the enzyme were heat treated at 65°C, which is the physiological temperature of *R. marinus*. Aggregation also occurred when the native protein was heated at 65°C. Analysis of the Cel12A sequence revealed that the protein, like certain other family 12 glycosyl hydrolases, comprises an amino terminal region that is highly hydrophobic (reminiscent of a signal peptide) and a  
15 catalytic domain which are linked by a linker moiety (e.g., region rich in proline, hydroxyamino acids (e.g., threonine, serine, tyrosine), alanine and/or glycine). Two variant polypeptides which lacked the amino-terminal hydrophobic region or the amino-terminal hydrophobic region and the linker moiety were produced in high yield when expressed in *E. coli*, indicating that cytotoxicity of Cel12A is mediated through  
20 the N-terminal hydrophobic region (putative signal peptide). The variant polypeptides were studied and their catalytic and physical properties compared to those of full-length Cel12A. The pH optima and substrate specificities of the variant polypeptides were the same as those of full-length Cel12A. However, the specific activities of the variants were about three fold higher than the specific activity of the full-length enzyme.  
25 Furthermore, the variant polypeptides were more stable than full-length Cel12A and had half-lives that exceeded 3 hours at 90°C and unfolding temperatures that were up to about 103°C.

### Polypeptides

- In one aspect, the invention relates to polypeptides that have thermostable cellulase activity. Preferably the polypeptides of the invention can hydrolyze  $\beta$ -1,4 bonds in  $\beta$ -1,4-linked and mixed linkage  $\beta$ -glucans (e.g., carboxymethyl cellulose, cellulose). The polypeptides of the invention have the amino acid sequence of a thermostable cellulase wherein one or more amino acids which are not part of the catalytic domain of the enzyme are deleted. The catalytic domain and other domains (e.g., signal peptide, linker domains) of a cellulase can be readily identified by analysis and comparison of the amino acid sequence of a thermostable cellulase with the sequences of other cellulases (see, for example, Henrissat, B., *Biochem J*, 280:309-316 (1991); Henrissat, B. *et al.*, *Biochem J*, 293:781-788 (1993)). For example, an amino acid sequence can be compared to the sequences in SWISS-PROT using a suitable comparison program, such as PROSITE (Hoffman, K. *et al.*, *Nucleic Acids Res*, 27:215-219 (1999)).
- In one example, the polypeptide of the invention can be a variant of a thermostable cellulase which contains an amino terminal hydrophobic region and a catalytic domain that are linked by a linker moiety (e.g., region rich in proline and/or hydroxylamino acids (e.g., threonine, serine, tyrosine)), such as certain family 12 glycosyl hydrolases (e.g., Cel12A from *R. marinus*, EglA from *Pyrococcus furiosus* and the like). The polypeptide of this example can have the amino acid sequence of the native or full-length cellulase (e.g., Cel12A, EglA) wherein one or more of the amino acids in the amino-terminal hydrophobic domain and/or linker moiety are deleted. Preferably, the polypeptide of this example has the amino acid sequence of the native or full-length cellulase wherein the amino acids which constitute the amino terminal hydrophobic region (e.g., signal peptide) or the amino acids which constitute the amino terminal hydrophobic region and the linker moiety are deleted. In certain embodiments, the deleted amino acid residues comprise a region in the range of at least the first 17 amino acid residues and up to the first 37 amino acid residues of the corresponding full-length polypeptide.

The nucleotide sequence of a nucleic acid encoding Cel12A of *Rhodothermus marinus* is deposited in GenBank under accession number U72637, the entire teaching of which are incorporated herein by reference. However, the originally deposited sequence contained sequence errors which have been rectified in the sequence depicted as (SEQ ID NO:1). Specifically, three nucleotides were missing, at positions 1019 (insert G), 1049 (insert C), 1093 (insert C), and positions 1467-68 CG should read GC and CG. The nucleic acid sequence (SEQ ID NO:1) has an open reading frame starting at position 710. The sequence of this open reading frame is presented herein as SEQ ID NO:3. The amino acid sequence of Cel12A of *Rhodothermus marinus* is presented herein as SEQ ID NO:2. The amino terminal hydrophobic domain of the protein (SEQ ID NO:2) consists of amino acid residues 1 to about 17, the amino acid residues from about position 18 to about position 37 constitute the linker moiety and the residues from about position 38 to position 261 constitute the catalytic domain. Note that due to the errors in the original DNA-sequence, the previously deposited amino acid sequence was erroneously assigned at residues 105-129.

In certain embodiments, the polypeptide of the invention has the amino acid sequence of Cel12A from *R. marinus* (SEQ ID NO:2) wherein one or more of the amino acid residues in the amino-terminal hydrophobic region and/or the linking moiety (positions 1 to about 40 of SEQ ID NO:2) are deleted. Preferably, the polypeptide of the invention has the amino acid sequence of SEQ ID NO:2 wherein at least about five or at least about ten amino acid residues in the amino-terminal hydrophobic region and/or the linking moiety are deleted. More preferably, the polypeptide of the invention has the amino acid sequence of SEQ ID NO:2 wherein at least about 15, or at least about 17, or at least about 20, or at least about 25, or at least about 30, or at least about 35 or about 37 of the amino acid residues in the amino-terminal hydrophobic region and/or the linking moiety are deleted. In additional embodiments, the polypeptide having thermostable cellulase activity has an amino acid sequence selected from the group consisting of residues 18-261 of SEQ ID NO:2, residues 19-261 of SEQ ID NO:2, residues 20-261 of SEQ ID NO:2, residues 21-261 of SEQ ID NO:2, residues 22-261 of

SEQ ID NO:2, residues 23-261 of SEQ ID NO:2, residues 24-261 of SEQ ID NO:2, residues 25-261 of SEQ ID NO:2, residues 26-261 of SEQ ID NO:2, residues 27-261 of SEQ ID NO:2, residues 28-261 of SEQ ID NO:2, residues 29-261 of SEQ ID NO:2, residues 30-261 of SEQ ID NO:2, residues 31-261 of SEQ ID NO:2, residues 32-261 of  
5 SEQ ID NO:2, residues 33-261 of SEQ ID NO:2, residues 34-261 of SEQ ID NO:2, residues 35-261 of SEQ ID NO:2, residues 36-261 of SEQ ID NO:2, residues 37-261 of SEQ ID NO:2 and residues 38-261 of SEQ ID NO:2. In more particular embodiments, the amino terminal amino acid residue of the polypeptide is methionyl. For example, the polypeptide can have the sequence of SEQ ID NO:2 wherein residues 2-17 or  
10 residues 2-37 are deleted. In another example, the amino acid sequence of the polypeptide is methionyl-residues 18-261 of SEQ ID NO:2 wherein the methionyl is the amino terminal residue of the polypeptide.

In a particularly preferred embodiment, the polypeptide having thermostable cellulase activity has the amino acid sequence of SEQ ID NO:2 wherein residues 2-17  
15 or residues 2-37 are deleted.

Further encompassed by the present invention are thermostable cellulase active polypeptides with substantial sequence identity to SEQ ID NO: 2, wherein one or more of the first 37 amino acid residues of the full-length sequence are deleted, such as e.g., the first 17 amino acids (the hydrophobic amino terminal region), and/or a portion of or  
20 all of the linker moiety comprising residues from about 18-37. Substantial sequence identity in this context is meant to include at least about 85% identity, preferably at least about 90% identity, such as about 95% or 97% identity, and also including about 99% sequence identity. "Sequence identity" as used herein is calculated based on a reference sequence, (which in this instance is the sequence of SEQ ID NO: 2). Algorithms for  
25 sequence analysis are known in the art, such as BLAST, described in Altschul *et al.*, *J. Mol. Biol.*, 215:403-10 (1990). Generally, the default settings with respect to e.g. "scoring matrix" and "gap penalty" will be used for alignment.

In additional embodiments, the polypeptide of the invention is a family 12 glycosyl hydrolase having the amino acid sequence of EglA from *Pyrococcus furiosus*



(SEQ ID NO:5) wherein one or more of the amino acid residues in the amino-terminal hydrophobic region and/or the linking moiety are deleted. The nucleotide sequence (SEQ ID NO:4) of a nucleic encoding EglA from *Pyrococcus furiosus* is deposited in GenBank under accession number AF181032, the entire teaching of which are  
5 incorporated herein by reference. The nucleic acid sequence (SEQ ID NO:4) has an open reading frame starting at position 82. The amino acid sequence of EglA from *Pyrococcus furiosus* is presented herein as SEQ ID NO:5. The amino terminal hydrophobic domain of EglA consists of amino acid residues 1 to about 19 and the amino acid residues from about position 28 to about position 49 constitute the linker  
10 moiety (Bauer, M.W. *et al.*, *J Bacteriology*, 181:284-290 (1999)). Accordingly, the polypeptide of the invention can comprise the amino acid sequence of residues 20 to 319 of SEQ ID NO:5 or the amino acid sequence of residues 50 to 319 of SEQ ID NO:5, for example.

Also encompassed by the present invention are thermostable cellulase active  
15 polypeptides with substantial sequence identity to SEQ ID NO: 5, wherein one or more of the first 49 amino acid residues (the hydrophobic amino terminal region and linker moiety) of the full-length sequence are deleted, such as e.g. the first 19 amino acids (the hydrophobic amino terminal region), and/or a portion of or all of the linker moiety comprising residues from about 20-49, where substantial sequence identity is defined as  
20 above.

The polypeptides of the invention can have improved characteristics, such as increased stability (e.g., thermal stability, detergent stability), increased solubility in aqueous solvents, increased catalytic activity (e.g., specific activity, catalytic rate) and/or reduced cytotoxicity relative to the native or full-length thermostable cellulase,  
25 but retain the substrate specificity of the native or full-length cellulase. Accordingly, the polypeptides of the invention provide many advantages over certain other thermostable cellulases. For example, as described herein large quantities of the polypeptides of the invention can be produced by expression in *E. coli*, or other host cells, including other bacterial cells, yeast cells or cells of a filamentous fungus. Examples of suitable

bacterial cells besides *E.coli*, are *Bacillus spp.*, *Lactobacillus spp.* and *Zymomomas spp.*; and of yeast cells are e.g., those of *Saccharomyces spp.* (particularly *S. cerevisiae*), *Schizosaccharomyces spp.*, and *Pichia spp.*; and useful cells of filamentous fungus include those from *Aspergillus spp.* such as *A. niger*, *A. nidulans* and *A. oryzae*; and

5 *Neurospora spp.*, e.g., *Neurospora crassa*.

In one embodiment, the polypeptide of the invention has enhanced stability (e.g, thermal stability, detergent stability, longer shelf life) relative to the corresponding full-length cellulase. For example, the catalytic activity of the polypeptide of the invention can have a half-life that is about 1.5, about two or more times longer than the

10 half-life of the full-length enzyme under certain conditions (e.g., storage at 4°C, heating to 90°C). In a particular embodiment, the catalytic activity of the polypeptide of the invention has a half-life of at least about 2.5 hours at 90°C. Preferably, the catalytic activity of the polypeptide of the invention has a half-life of at least about 3 hours, or about 5 hours, or about 10 hours or more at 90°C.

15 In another embodiment, the polypeptide of the invention has enhanced specific activity relative to the corresponding full-length cellulase. For example, the specific activity of the polypeptide of the invention can be about 1.5 or about two or more times greater than the specific activity of the corresponding full-length cellulase. In a particular embodiment, the specific activity of the polypeptide of the invention is at least

20 about two times greater than the specific activity of the corresponding full-length cellulase. Preferably, the specific activity of the polypeptide of the invention is at least about 3, or about 5 or more times greater than the specific activity of the corresponding full-length cellulase. The specific activity of the polypeptide of the invention and of full-length cellulases can be determined using any suitable method, such as the method

25 described herein. Preferably, the specific activity is determined using carboxymethyl cellulose as the substrate.

The polypeptide of the invention can be a fusion protein comprising a polypeptide having thermostable cellulase activity as described herein which is fused to a second moiety not occurring in the cellulase as found in nature. Thus, the second

moiety can be an amino acid, oligopeptide or polypeptide. The second moiety can be linked to the first moiety at a suitable position, for example, the N-terminus, the C-terminus or internally. In one embodiment, the fusion protein comprises an affinity ligand (e.g., an enzyme, an antigen, epitope tag (e.g., hemagglutinin (HA), HSV-Tag (SQPELAPEDPED (SEQ ID NO:6)), polyhistidine (e.g., His<sub>6</sub>), a binding domain) as the first moiety, and a second moiety comprising a linker sequence and polypeptide that has thermostable cellulase activity as described herein. Additional (e.g., third, fourth) moieties can be present as appropriate. When the polypeptide is produced as a fusion protein, the fusion partner (e.g., HA, HSV-Tag, His<sub>6</sub>) can be used to facilitate purification and/or isolation. If desired, the fusion partner can then be removed from polypeptide of the invention (e.g., by proteolytic cleavage).

#### *Nucleic acids and Constructs*

The invention also relates to isolated nucleic acids and to constructs comprising the nucleic acids. The nucleic acids of the invention can be DNA or RNA, for example, mRNA. The nucleic acid molecules can be double-stranded or single-stranded; single stranded RNA or DNA can be either the coding, or sense, strand or the non-coding, or antisense, strand. Preferably, the nucleic acids encode a polypeptide of the invention. If desired, the nucleotide sequence of the isolated nucleic acid can include additional non-coding sequences such as non-coding 3' and 5' sequences (including regulatory sequences, for example). Additionally, the nucleic acids of the invention can be fused to a nucleic acid comprising a marker sequence, for example, a nucleotide sequence which encodes a polypeptide to assist in isolation or purification of the polypeptide. Representative sequences include, but are not limited to, those which encode a glutathione-S-transferase (GST) fusion protein, a poly-histidine (e.g., His<sub>6</sub>), hemagglutinin, HSV-Tag, for example.

The nucleic acid molecules of the invention are "isolated" as used herein, an "isolated" nucleic acid molecule or nucleotide sequence is intended to mean a nucleic acid molecule or nucleotide sequence which is not flanked by nucleotide sequences

which normally flank the gene or nucleotide sequence (as in genomic sequences) and/or has been completely or partially purified from other nucleic acids (e.g., as in an DNA or RNA library). For example, an isolated nucleic acid of the invention may be substantially isolated with respect to the complex cellular milieu in which it naturally occurs. In some instances, the isolated material will form part of a composition (for example, a crude extract containing other substances), buffer system or reagent mix. In other circumstance, the material may be purified to essential homogeneity, for example as determined by PAGE or column chromatography such as HPLC. Thus, an isolated nucleic acid molecule or nucleotide sequence can include a nucleic acid molecule or nucleotide sequence which is synthesized chemically, using recombinant DNA technology or using any other suitable method. Therefore, a recombinant nucleic acid (e.g., DNA, RNA) contained in a vector is included in the definition of "isolated" as used herein. Also, isolated nucleotide sequences include recombinant nucleic acid molecules (e.g., DNA, RNA) in heterologous organisms, as well as partially or substantially purified nucleic acids in solution. *In vivo* and *in vitro* RNA transcripts of a DNA molecule of the present invention are also encompassed by "isolated" nucleotide sequences.

The present invention also pertains to nucleotide sequences which are not necessarily found in nature but which encode a polypeptide of the invention. For example, DNA molecules which comprise a sequence which is different from the naturally-occurring nucleotide sequence (e.g., SEQ ID NO:3) but which, due to the degeneracy of the genetic code, encode a polypeptides of the invention are the subject of this invention. The invention also encompasses variations of the nucleotide sequences of the invention, such as those encoding active fragments or active derivatives of the polypeptides as described below. Such variations can be naturally-occurring, or non-naturally-occurring, such as those induced by various mutagens and mutagenic processes. Intended variations include, but are not limited to, addition, deletion and substitution of one or more nucleotides which can result in conservative or non-conservative amino acid changes, including additions and deletions. Preferably, the

nucleotide or amino acid variations that are within the catalytic domain are silent or conserved; that is, they do not alter the improved characteristics or activity of the encoded polypeptide.

The invention described herein also relates to fragments of the isolated nucleic acid molecules described herein. The term "fragment" is intended to encompass a portion of a nucleotide sequence described herein which is from at least about 25 contiguous nucleotides to at least about 50 contiguous nucleotides or longer in length; such fragments are useful as probes and also as primers. Particularly preferred primers and probes selectively hybridize to the nucleic acid molecule encoding the polypeptides described herein. For example, fragments which encode polypeptides that retain activity, as described below, are particularly useful.

The invention also pertains to nucleic acid molecules which hybridize under high stringency hybridization conditions, such as for selective hybridization, to a nucleotide sequence described herein (e.g., nucleic acid molecules which specifically hybridize to a nucleotide sequence encoding polypeptides described herein and encode a polypeptide having thermostable cellulase activity). Hybridization probes include synthetic oligonucleotides which bind in a base-specific manner to a complementary strand of nucleic acid. Suitable probes include polypeptide nucleic acids, as described in Nielsen *et al.*, *Science*, 254:1497-1500 (1991).

Such nucleic acid molecules can be detected and/or isolated by specific hybridization (e.g., under high stringency conditions). "Stringency conditions" for hybridization is a term of art which refers to the incubation and wash conditions, e.g., conditions of temperature and buffer concentration, which permit hybridization of a particular nucleic acid to a second nucleic acid; the first nucleic acid may be perfectly (i.e., 100%) complementary to the second, or the first and second may share some degree of complementarity which is less than perfect (e.g., 60%, 75%, 85%, 95%). For example, certain high stringency conditions can be used which distinguish perfectly complementary nucleic acids from those of less complementarity.

"High stringency conditions", "moderate stringency conditions" and "low stringency conditions" for nucleic acid hybridizations are explained on pages 2.10.1-2.10.16 and pages 6.3.1-6 in *Current Protocols in Molecular Biology* (Ausubel, F.M. *et al.*, "*Current Protocols in Molecular Biology*", John Wiley & Sons, (1998)) the teachings of which are hereby incorporated by reference. The exact conditions which determine the stringency of hybridization depend not only on ionic strength (e.g., 0.2XSSC, 0.1XSSC), temperature (e.g., room temperature, 42°C, 68°C) and the concentration of destabilizing agents such as formamide or denaturing agents such as SDS, but also on factors such as the length of the nucleic acid sequence, base composition, percent mismatch between hybridizing sequences and the frequency of occurrence of subsets of that sequence within other non-identical sequences. Thus, high, moderate or low stringency conditions can be determined empirically.

By varying hybridization conditions from a level of stringency at which no hybridization occurs to a level at which hybridization is first observed, conditions which will allow a given sequence to hybridize (e.g., selectively) with the most similar sequences in the sample can be determined.

Exemplary conditions are described in Krause, M.H. and S.A. Aaronson, *Methods in Enzymology*, 200:546-556 (1991). Also, in, Ausubel, *et al.*, "*Current Protocols in Molecular Biology*", John Wiley & Sons, (1998), which describes the determination of washing conditions for moderate or low stringency conditions. Washing is the step in which conditions are usually set so as to determine a minimum level of complementarity of the hybrids. Generally, starting from the lowest temperature at which only homologous hybridization occurs, each degree (°C) by which the final wash temperature is reduced (holding SSC concentration constant) allows an increase by 1% in the maximum extent of mismatching among the sequences that hybridize. Generally, doubling the concentration of SSC results in an increase in  $T_m$  of -17EC. Using these guidelines, the washing temperature can be determined empirically for high, moderate or low stringency, depending on the level of mismatch sought.

For example, a low stringency wash can comprise washing in a solution containing 0.2XSSC/0.1% SDS for 10 min at room temperature; a moderate stringency wash can comprise washing in a prewarmed solution (42°C) solution containing 0.2XSSC/0.1% SDS for 15 min at 42°C; and a high stringency wash can comprise washing in prewarmed (68°C) solution containing 0.1XSSC/0.1%SDS for 15 min at 68°C. Furthermore, washes can be performed repeatedly or sequentially to obtain a desired result as known in the art.

Equivalent conditions can be determined by varying one or more of the parameters given as an example, as known in the art, while maintaining a similar degree of identity or similarity between the target nucleic acid molecule and the primer or probe used. Hybridizable nucleotide sequences are useful as probes and primers for identification of organisms (e.g., recombinant bacteria) containing a nucleic acid of the invention and to isolate a nucleic acid of the invention, for example. As used herein, the term "primer" refers to a single-stranded oligonucleotide (e.g., synthetic oligo deoxyribonucleotide) which acts as a point of initiation of template-directed DNA synthesis under appropriate conditions (e.g., in the presence of four different nucleoside triphosphates and an agent for polymerization, such as, DNA or RNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature. The appropriate length of a primer depends on the intended use of the primer, but typically ranges from 15 to 30 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template, but must be sufficiently complementary to hybridize with a template. The term "primer site" refers to the area of the target DNA to which a primer hybridizes. The term "primer pair" refers to a set of primers including a 5' (upstream) primer that hybridizes with the 5' end of the DNA sequence to be amplified and a 3' (downstream) primer that hybridizes with the complement of the 3' end of the sequence to be amplified.

The invention also pertains to nucleotide sequences which have substantial identity with the nucleotide sequences described herein; particularly preferred are

nucleotide sequences which have at least about 10%, preferably at least about 20%, more preferably at least about 30%, more preferably at least about 40%, even more preferably at least about 50%, yet more preferably at least about 70%, still more preferably at least about 80%, and even more preferably at least about 90% or 95% identity, with nucleotide sequences described herein. Particularly preferred in this instance are nucleotide sequences encoding polypeptides having thermostable cellulase activity and an amino acid sequence described herein.

To determine the percent identity of two nucleotide sequences, the sequences can be aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first nucleotide sequence). The nucleotides at corresponding nucleotide positions can then be compared. When a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical positions/total # of positions x 100).

The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:5873-5877 (1993). Such an algorithm is incorporated into the NBLAST program which can be used to identify sequences having the desired identity to nucleotide sequences of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, *Nucleic Acids Res*, 25:3389-3402 (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., NBLAST) can be used. See ~~<http://www.ncbi.nlm.nih.gov>~~. In one embodiment, parameters for sequence comparison can be set at W=12. Parameters can also be varied (e.g., W=5 or W=20). The value "W" determines how many continuous nucleotides must be identical for the program to identify two sequences as containing regions of identity.

mm  
5/14/03



In certain embodiments, the isolated nucleic acid encodes a polypeptide having the amino acid sequence of Cel12A from *R. marinus* (SEQ ID NO:2) wherein one or more of the amino acid residues in the amino-terminal hydrophobic region and/or the linking moiety (positions 1 to about 37 of SEQ ID NO:2) are deleted. Preferably, the isolated nucleic acid encodes a polypeptide having the amino acid sequence of SEQ ID NO:2 wherein at least about five or at least about ten amino acid residues in hydrophobic region and/or the linking moiety are deleted. More preferably, the isolated nucleic acid encodes a polypeptide having the amino acid sequence of SEQ ID NO:2 wherein at least about 15, or at least about 17, or at least about 20, or at least about 25, or at least about 30, or at least about 35 or about 37 of the amino acid residues in hydrophobic region and/or the linking moiety are deleted. In additional embodiments, the isolated nucleic acid of the invention encodes a polypeptide having thermostable cellulase activity, and has a nucleotide sequence selected from the group consisting of nucleotides 52-783 of SEQ ID NO:3, nucleotide 55-783 of SEQ ID NO:3, nucleotides 58-783 of SEQ ID NO:3, nucleotides 61-783 of SEQ ID NO:3, nucleotides 64-783 of SEQ ID NO:3, nucleotides 67-783 of SEQ ID NO:3, nucleotides 70-783 of SEQ ID NO:3, nucleotides 73-783 of SEQ ID NO:3, nucleotides 76-783 of SEQ ID NO:3, nucleotides 79-783 of SEQ ID NO:3, nucleotides 82-783 of SEQ ID NO:3, nucleotides 85-783 of SEQ ID NO:3, nucleotides 88-783 of SEQ ID NO:3, nucleotides 91-783 of SEQ ID NO:3, nucleotides 94-783 of SEQ ID NO:3, nucleotides 97-783 of SEQ ID NO:3, nucleotides 100-783 of SEQ ID NO:3, nucleotides 103-783 of SEQ ID NO:3, nucleotides 106-783 of SEQ ID NO:3, nucleotides 109-783 of SEQ ID NO:3 and nucleotides 112-783 of SEQ ID NO:3. In more particular embodiments, the amino terminal amino acid residue of the polypeptide is methionyl. Accordingly, the isolated nucleic acid of the invention can have the sequence atg-nucleotides 52-783 of SEQ ID NO:3 or atg-nucleotides 112-783 of SEQ ID NO:3, for example. In a particularly preferred embodiment, the isolated nucleic acid encoding a polypeptide having thermostable cellulase activity has the sequence of nucleotides 52-789 of SEQ ID NO:3 or nucleotides 112-783 of SEQ ID NO:3.

In additional embodiments, the isolated nucleic acid of the invention encodes a family 12 glycosyl hydrolase having the amino acid sequence of EglA from *Pyrococcus furiosus* wherein one or more of the amino acid residues in the amino-terminal hydrophobic region and/or the linking moiety are deleted. For example, the isolated  
5 nucleic acid can have the sequence of nucleotides 139-1041 of SEQ ID NO:4 or the sequence of nucleotides 229-1041 of SEQ ID NO:4.

The nucleic acids described herein can be amplified by methods known in the art. For example, amplification can be accomplished by the polymerase chain reaction (PCR). See generally *PCR Technology: Principles and Applications for DNA*  
10 *Amplification* (ed. H.A. Erlich, Freeman Press, NY, NY, 1992); *PCR Protocols: A Guide to Methods and Applications* (eds. Innis, et al., Academic Press, San Diego, CA, 1990); Mattila *et al.*, *Nucleic Acids Res.* 19:4967 (1991); Eckert *et al.*, *PCR Methods and Applications* 1, 17 (1991); PCR (eds. McPherson *et al.*, IRL Press, Oxford); and U.S. Patent 4,683,202. Other suitable amplification methods include the ligase chain  
15 reaction (LCR) (see Wu and Wallace, *Genomics*, 4:560 (1989), Landegren *et al.*, *Science*, 241:1077 (1988), transcription amplification (Kwoh *et al.*, *Proc. Natl. Acad. Sci. USA*, 86:1173 (1989)), and self-sustained sequence replication (Guatelli *et al.*, *Proc. Nat. Acad. Sci. USA*, 87:1874 (1990)) and nucleic acid based sequence amplification (NASBA).

20 The invention includes recombinant constructs comprising an isolated nucleic acid as described herein operably linked to at least one regulatory sequence. Generally, the isolated nucleic acid is joined (e.g., ligated, inserted) to a suitable vector, such as a plasmid or viral vector that includes an origin of replication, a selectable genetic marker and/or sequences which allow the insert nucleic acid to be expressed (transcribed  
25 and/or translated). For example the recombinant construct can be an expression vector containing a nucleic acid sequence encoding a polypeptide described herein that is operably linked to at least one regulatory sequence. Many expression vectors are commercially available, and other suitable vectors can be readily prepared by the skilled artisan. As used herein, "operably linked" means that the nucleotide sequence is linked

to a regulatory sequence in a manner which allows expression (e.g., transcription and/or translation) of the nucleic acid sequence. Regulatory sequences are art-recognized and are selected to produce the polypeptide or active derivative or fragment thereof. Accordingly, the term "regulatory sequence" includes promoters, enhancers, ribosome binding sites, polyadenylation signals and other expression control elements which are described, for example, in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). For example, regulatory sequences native to *R. marinus* can be employed to express Cel12A. Suitable expression vectors and regulatory elements can be selected to provide for the desired level of expression in a desired host cell. For example, the polypeptides of the present invention can be produced by ligating a nucleic acid encoding the polypeptide into a vector suitable for expression in an appropriate host cell (see, for example, Broach, *et al.*, *Experimental Manipulation of Gene Expression*, ed. M. Inouye (Academic Press, 1983) p. 83; *Molecular Cloning: A Laboratory Manual*, 2nd Ed., ed. Sambrook *et al.* (Cold Spring Harbor Laboratory Press, 1989) Chapters 16 and 17)). For example, where expression in *E. coli* is desired the vector can include the T7/lac promotor. Other suitable regulatory sequences for use in *E. coli*, other bacteria, fungi (yeast, molds), insect and/or mammalian cells are known in the art. Typically, expression constructs contain one or more suitable selectable markers. Suitable selectable genetic markers include, for example, genes which confer resistance to antibiotics such as the  $\beta$ -lactamase gene from Tn3, the kanamycin-resistance gene from Tn903, the chloramphenicol-resistance gene from Tn<sub>9</sub>, and the like. Suitable origins or replication include, for example, Ori p, colE1 Ori and the like.

#### *Recombinant Host Cells and Methods of Production*

The invention also relates to recombinant host cells (e.g., prokaryotic, eukaryotic, archaeon) which comprise an expression vector or isolated nucleic acid as described herein. For example, cells which can comprise a vector or isolated nucleic of the invention (for example, as a result of transformation, transfection or transduction)

include, but are not limited to, bacterial cells (e.g., *R. marinus*, *E. coli* (e.g., *E. coli* K12 strains), *Streptomyces*, *Pseudomonas*, *Bacillus*, *Serratia marcescens*, *Salmonella typhimurium*), fungi including yeasts (e.g., *Saccharomyces cerevisiae*, *Pichia pastoris*) and molds (e.g., *Aspergillus sp.*), insect cells (e.g., Sf9) or mammalian cells (e.g., COS, CHO). The construct (e.g., expression vector) can be introduced into the host cells using any suitable method (e.g., electroporation, transfection using calcium chloride, rubidium chloride, calcium phosphate, DEAE-dextran, or other substances; microprojectile bombardment; lipofection, infection, transduction).

The isolated nucleic acid molecules and vectors of the invention are useful in the manufacture of the encoded polypeptide, as probes for isolating homologous sequences (e.g., from other bacteriophage species), as well as for detecting the presence of a nucleic acid of the invention in a cells.

The invention also relates to a method of producing (*in vitro*, *in vivo*) a polypeptide having thermostable cellulase activity. For example, a nucleic acid encoding a polypeptide of the invention, or a construct comprising such nucleic acid, can be introduced into a suitable host cell by a method appropriate to the host cell selected (e.g., transformation, transfection, electroporation, infection), such that the nucleic acid is operably linked to one or more expression control elements (e.g., in a vector, in a construct created by processes in the cell, integrated into the host cell genome). Host cells can be maintained under conditions suitable for expression (e.g., *in vivo* or *in vitro*, in the presence of inducer, suitable media supplemented with appropriate salts, growth factors, antibiotic, nutritional supplements, etc.), whereby the encoded polypeptide is produced. In additional embodiments, the polypeptide of the invention can be produced by *in vitro* translation of a nucleic acid that encodes the polypeptide (e.g., a nucleic acid having a sequence described herein), by chemical synthesis (e.g., solid phase peptide synthesis) or by any other suitable method. If desired, the polypeptide having thermostable cellulase activity can be isolated (e.g., from the host cells, medium, milk). It will be appreciated that the method encompasses

expression of the polypeptide in a host cell of a transgenic animal or plant (see e.g., U.S. Patent Nos. 6,013,857, 5,990,385, 5,994,616).

*Uses for Polypeptides that have Thermostable Cellulase Activity*

The polypeptides of the invention are useful in a variety of applications, such as industrial processes. For example, the polypeptides can be used to improve freeness and to remove inks, coatings, toners and colors from wood or paper pulp (see, e.g., U.S. Patent Nos. 6,066,233, 5,525,193 and WO 98/44189). The polypeptides of the invention can also be included in detergent compositions and to treat cellulose-containing textiles and garments to improve the feel of the fabric or to remove lint, for example (see, e.g., U.S. Patent Nos. 5,904,736, 5,792,641, 5,445,957). Additional uses for the polypeptides of the invention are in the treatment of fibrous crops, fruits and vegetables or grains to improve feed values or to extract starches (e.g., sugars) or other components of the crop, fruit, vegetable or grain being treated. A number of methods for treating crops, fruits, vegetables or grains with a cellulase are known in the art (see, e.g., U.S. Patent Nos. 5,948,454, 4,795,101, 4,371,552 and WO 97/23652).

EXAMPLE

MATERIALS AND METHODS

*Bacterial strains*

Nova Blue and BL21 (DE3) strains of *E. coli* (Novagen, Madison, WI) were used for cloning and expression of variants of Cel12A. The GE2205 (Institute of Biology, University of Iceland) strain of *E. coli* containing plasmid pET23bAH was used for expression of a *cel12A* gene (Halldorsdottir, S. *et al.*, *Appl Microbiol Biotechnol*, 49:277-284 (1998)). Plasmid pET25b(+) (Novagen) was used for expression of variants of the Cel12A protein.

*Construction of plasmids and sequencing*

A gene fragment encoding 223 amino acids of the Cel12A (starting at the 38th amino acid), named  $\delta$  cel12A with introduced *NdeI* and *HindIII* restriction sites was obtained by PCR utilising pET23bAH plasmid (Halldorsdottir, S. *et al.*, *Appl Microbiol Biotechnol*, 49:277-284 (1998)) as a template. The following primers were used for amplification of  $\delta$  cel12A: forward synthetic oligonucleotide primer (*NdeI* site underlined): 5'-AGGACTCCCATATGACCGTCGAGCTGTCGG-3' (SEQ ID NO:7), reverse synthetic oligonucleotide primer (*HindIII* site underlined): 5'-ACCTGAGAAAGCTTCTGCACCGTTACGGA-3' (SEQ ID NO:8). AmpliTaq Gold™ (Perkin-Elmer AB, Foster City, CA) DNA polymerase and hot start PCR protocol of 35 cycles of denaturation at 95°C for 1 min, annealing at 60°C for 1 min and primer extension at 72°C for 1 min were used. PCR fragments were separated by agarose gel electrophoresis, purified using a QIAEX II kit (QIAGEN GmbH, Hilden, Germany), digested overnight with *Nde I* and *Hind III* and purified again using the QIAEX II kit.

The gene fragment encoding Cel12A with the amino terminal hydrophobic region (putative signal peptide (first 17 amino acids)) deleted, named *cel12A(sp-)* was amplified using the following primers: forward synthetic oligonucleotide primer (*NdeI* site underlined): 5'-AGCTCCCATATGTGCGACTGGCTCTTTCC-3' (SEQ ID NO:9) and reverse synthetic oligonucleotide primer (*HindIII* site underlined): 5'-ACCTGAGAAAGCTTCTGCACCGTTACGGA-3' (SEQ ID NO:10) and purified as described above. The pET25b(+) vector was opened with restriction enzymes (*Nde I* and *Hind III*) and purified using QIAEX II kit. Inserts were cloned into appropriately opened vectors to produce plasmids: pET25 $\delta$ Cel12AH and pET25Cel12A(sp-)H.

Plasmids were prepared using QIAGEN Mini Plasmid Kit. The presence of cloned inserts in the plasmids was confirmed by restriction analysis and partial sequencing by the dideoxy chain termination method using the ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction (Perkin-Elmer).

All restriction enzymes and T4 ligase were from Life Technologies (Rockville, MD). The enzymes were used according to the manufacture's protocols.

#### *Culture conditions and protein production*

For expression of variant forms of the cellulase, bacteria were grown in baffled  
5 flasks in LB medium containing ampicilin to  $OD_{600} = 0.7$  and then induced with IPTG  
(final concentration 1mM). Growth was continued for an additional 2.5 - 3 hours. Cells  
were harvested by centrifugation. The cell pellet was resuspended in 20 mM Tris-HCl  
pH 7.4 / 20 mM imidazole / 0.5 M NaCl buffer. Disintegration of the cells was  
performed using a UP400S sonicator (Dr. Heilscher GmbH, Stahnsdorf, Germany) at 60  
10 W  $cm^{-2}$ , 3 x 120 seconds on ice. The crude cell extract was centrifuged at 27,000 x g for  
20 min at 4°C and the supernatant stored frozen in -20°C for later purification.

Fed-batch cultivation for production of the Cel12A protein was carried out as  
described (Nordberg, K.E. *et al.*, *J. Ferment. Bioeng.*, (in press, 2000)). Bacteria were  
grown until the  $OD_{600}$  reached 32 and induced by addition of IPTG (final concentration  
15 0.5 mM). Growth was continued for an additional 90 min. Whole fermentation culture  
was run three times on a Gaulin high-pressure homogenizer (APV-Schröder GmbH,  
Lübeck, Germany) at 700 atm to disintegrate the cells. The cell extract was centrifuged  
(27,000 x g, 4 °C, 15 min) and the supernatant frozen at -20°C and stored for  
purification.

#### 20 *Protein purification*

Crude cell extracts containing variant Cel12A proteins were heat-treated at 65°C  
for 30 min, centrifuged at 27,000 x g for 15 min and the supernatants were loaded onto a  
chromatography column (crude extracts containing full-length Cel12A were not  
heat-treated). Recombinant proteins were purified utilizing C-terminal 6xHisidine tags  
25 and immobilized metal-ion affinity chromatography (IMAC) as described  
(Abou-Hachem *et al.*, *Biochem. J.*, 345:53-60 (2000)). The eluate was dialyzed three  
times for several hours, against 45 volumes of 20 mM Tris-HCl pH 7.5 in dialysis tubes

with a molecular weight cut-off of 3.5 kD. Proteins were concentrated using Aquacide III (CALBIOCHEM, La Jolla, CA) according to the manufacture's protocol. Then, the concentrated protein solutions were centrifuged and the supernatants were stored at 4°C until used for further testing.

5           The histidine tag was removed from the purified polypeptides by digestion with carboxypeptidase-A type II from bovine pancreas (Sigma) for 1 hour at 37°C. Following digestion, the polypeptide solutions were loaded onto the affinity column and portions which did not bind (flow through) were collected. A portion of the flow through was analysed by SDS-PAGE to verify that the histidine tag had been removed.

10           Gel filtration chromatography was used to separate of cleaved and uncleaved Cell12A(sp-)H protein. A 50 cm column with internal diameter of 17 mm (Biorad), packed with Sephadex G-50 resin (Amersham Pharmacia Biotech AB, Uppsala, Sweden) was used. The column was run using 20 mM Tris-HCl pH 7.5.

#### *Protein characterization*

15           Protein production was analysed by sodium dodecyl sulphate polyacrylamide electrophoresis (SDS-PAGE) in 12.5 % gels using the method of Laemmli (Laemmli, U.K., *Nature*, 227:680-685 (1970)). Proteins were visualized by staining with Coomassie brilliant blue G250 (Merck, Darmstadt, Germany). Enzyme activity was detected using a 1 % (w/v) agarose overlay gel containing 0.02 % (w/v)

20   carboxymethyl cellulose (CMC). The SDS/PAGE gels were washed with phosphate buffer pH 7.0 containing 1% Triton X-100 (t-octylphenoxypolyethoxyethanol, Merck) for 20 min. Thereafter, the gels were washed with phosphate buffer pH 7.0, covered with the overlay gel and incubated for 2 hours at 65°C. Then, the agarose gels were stained in 1 % (w/v) Congo red solution. Excess dye was removed by washing with 1

25   M NaCl.

The dinitrosalicylic acid (DNS) stopping method was used for enzyme activity measurements. Samples were incubated with 0.9 % (w/v) CMC in 20 mM Tris-HCl pH 7.5 at 65°C for about 20 minutes or about 30 minutes. The enzymatic reactions were



stopped by adding 1.5 volume of DNS-solution (1% DNS / 0.2% phenol / 0.05% Na<sub>2</sub>SO<sub>3</sub> / 1% NaOH / 20% Na-K-tartrate) to the samples and boiling the samples for 15 minutes. The amount of reducing sugars released was measured in 96 well microtiter plates by absorbance at 550 nm, using glucose solutions as standards.

- 5 CMC, phosphoric acid swollen cellulose (PASC), birch wood xylan, lichenan, locust bean gum and guar gum (all SIGMA, St. Louis, MO) and Konjak (Megazyme International Ltd.) were used for enzyme substrate specificity tests. Temperature optima were determined by running the activity assay at different temperatures ranging from 40°C to 105°C in 20 mM Tris-HCl pH 7.5 for 20 minutes. pH optima were studied by
- 10 running the activity assay in 0.1 M citrate/phosphate buffer (pH ranging from 3.0 to 6.0), 0.1 M sodium phosphatebuffer (pH ranging from 6.0 to 7.0), 50mM Tris-HCl buffer (pH ranging from 7.0 to 9.0) and glycine-NaOH buffer (pH ranging from 9.0 to 10.0) at 65°C for 30 min. Temperature stability was examined by determining residual enzymatic activity after incubating the proteins in absence of substrates at 65°C, 85°C
- 15 and 90°C for predetermined periods of time.

#### *DNA and protein sequence analysis*

- A few Internet data banks and net tools were utilized for primary sequence analysis and secondary and tertiary structure prediction. BLAST 2.0 (Altschul *et al.*, *Nucleic Acids Research*, 25:3389-3402 (1997); Altschul *et al.*, *J Mol Bio*, 215:403-410
- 20 (1990)) which is available on a server of the National Center for Biotechnology Information - <http://www.ncbi.nlm.nih.gov/>) was used to investigate related sequences. Primary structure analysis was done using tools and databases available on the ExPASy Molecular Biology Server (<http://www.expasy.ch/>), such as PROSITE (Hofmann *et al.*, *Nucleic Acids Research*, 27:215-219 (1999); Bairoch A., *Nucleic Acids Research* 20
  - 25 *Suppl*:2013-2018 (1992)) and SignalP (Nielsen H. *et al.*, *Protein Engineering*, 10:1-6 (1997)).

### *Differential scanning calorimetry (DSC)*

Thermal transitions were measured by DSC using a VP-DSC microcalorimeter (MicroCal Inc., Northampton, MA), and the following parameters: scan rate 60°C/h and temperature range from 25 °C to 130 °C. Different protein concentrations (ranging from 5 0.1 to 0.4 mg/ml) were used and measurements were performed in 20 mM Tris-HCl pH 7.5. Additional measurements were made using the same Tris buffer supplemented with 1mM CaCl<sub>2</sub>, 0.5 % CMC or 0.5 % SDS.

## RESULTS

### *Expression of full length Cel12A in E. coli*

10 Cultivation of *E. coli* transformed with an expression plasmid encoding full length Cel12A in a shake-flask, revealed that induction of expression of the recombinant protein resulted in a decrease in optical density measured at 600 nm (Fig. 1), indicating that full length Cel12A induced cell lysis. Significant changes in bacterial morphology were noted after induction of Cel12A expression. Notably, the cultures 15 contained mostly long (up to 30 fold longer than normal *E. coli* cell) filamentous bacteria and multiple-cell clumps. Production of the cellulase (i.e., Cel12A) was extremely low due to the toxicity of the full length protein, and no clear recombinant protein band was visible on SDS-polyacrylamide gels (Fig. 2). However, two faint bands were detected on an activity stained over-layer gel. The second band detected 20 using the activity over-layer gel was probably produced by cleavage of Cel12A within linker region by bacterial proteases. Similar results were obtained when full length Cel12A was expressed in *E. coli* at 30°C. Specifically, no Cel12A was detected by SDS-PAGE, however no significant decrease in optical density was noticed.

To obtain enough protein for further tests the protein was produced using 25 fed-batch cultivation. However, induction of expression at high cell density led to a decrease in optical density that was associated with intensive foaming and increased culture viscosity, hallmarks of cell lysis. As in the shake-flask cultures, outgrowth of bacteria was eventually observed in cultures maintained under growth conditions. The

cultivation was stopped 2.5 hours after induction. Enzymatic activities (measured with DNS-stopping method) found in the cell pellet and in the supernatant were about equal. Thus, the Cel12A protein was purified from the entire batch culture (cells and media) to avoid protein lost. The Cel12A was purified by IMAC chromatography without prior  
 5 heat-treatment of crude cell extract. Partially purified protein was then heat-treated. Intensive precipitation of recombinant protein occurred upon heating, resulting in very low yield of pure Cel12A.

The purified full length Cel12A protein was not stable at 4°C and lost activity after several weeks of storage.

#### 10 *Sequence analysis of the cel12A gene*

Properties of the Cel12A based on sequence analysis were predicted using ProtScale (<http://www.expasy.ch/cgi-bin/protscale.pl>). Amino acid sequence profiling using an optimised matching hydrophobicity (OMH) scale (Sweet, R.M. *et al.*, *J Mol Biol*, 171:479-488 (1983)) demonstrated that the N-terminal part of Cel12A is the most  
 15 hydrophobic region of the molecule. This hydrophobic region is connected to a short and less hydrophobic region. Profiling using the average flexibility index (Bhaskaran, R. *et al.*, *Int J Pept Protein Res*, 32:242-255 (1988)) showed that the first several amino acids of the protein (amino terminal amino acids) could form highly rigid secondary structure in the protein. Additionally, the hydrophobic amino terminal region was  
 20 followed by the most flexible region of the protein. The flexible region comprises a doublet of glutamic acid and proline that is repeated four times ([EP]<sub>4</sub>). These studies revealed that Cel12A includes a catalytic domain and a hydrophobic N-terminal region that are linked through a flexible linker moiety (Fig. 3A). The amino-terminal region (first 38 amino acids) was shown to be toxic when expressed in BL21(DE3) bacteria  
 25 (Fig. 1).

### *Expression of Cel12A variants*

To overcome bacterial host death during expression of the Cel12A, variant proteins were designed. (See schematics of proteins presented in Figs. 3A-3D.) A gene fragment encoding the catalytic domain ( $\delta$ Cel12AH) downstream from the [EP]<sub>4</sub> linker  
 5 was cloned under the control of T7/lac promoter in plasmid pET25b and expressed in *E. coli* BL21(DE3). The variant protein accumulated in cytoplasm of the host bacteria and no bacterial lysis was observed (Fig. 1). The growth rate of the bacteria was slightly inhibited due to the intensive production of the recombinant variant protein which accounted for about 40 % of total protein content of the cells (Fig. 2). The variant  
 10 protein,  $\delta$ Cel12A, did not precipitate when incubated at 65°C and retained activity on CMC as assessed by overlay gel test.

A DNA fragment encoding a variant Cel12A that lacked the signal peptide (first 17 amino acids) was cloned into the pET25b plasmid and expressed in *E. coli*. Bacteria effectively produced active protein without cell lysis. However, the recombinant variant  
 15 cellulase, Cel12A(sp-)H, was partially cleaved by bacterial protease and two bands were observed on SDS-polyacrylamide gels (Fig. 2), as well as on activity gels. Both variant cellulases were stable, and were stored at 4°C for several weeks.

### *Characterization of Cel12A and variants*

The full-length cellulase expressed in *E. coli* has been characterised  
 20 (Halldorsdottir, S. *et al.*, *Appl Microbiol Biotechnol*, 49:277-284 (1998)). However, all activity tests described herein were conducted using full-length Cel12A and all variants. The full-length enzyme and all variants were rested with and without the His-tag (His<sub>6</sub>), and no differences in properties were observed. The effect of temperature on activity was investigated by incubating the proteins at temperatures ranging from 40°C to 105°C  
 25 and then assaying for enzymatic activity. The activity of full-length Cel12A and Cel12A(sp-)H increased up to 105°C, while Cel12AH displayed maximal activity at about 90°C (Table 1). The variant forms of the cellulase retained 95% - 100% of their initial activity after 16 hours at 85°C. The half-life of activity at 90°C for full-length

Cell12A was 3 hours, for Cell12AH was 2 hours and for Cell12(sp-)H was 5 hours. The pH optima for the full-length enzyme and the variants were similar. All proteins were active over a broad pH range and expressed over 40% of maximal activity at pH ranging from about 4.0 to about 8.0 and over 20% of maximal activity at pH 10.

- 5           The variant Cell12A proteins had the same substrate specificities as full-length Cell12A protein and hydrolysed soluble polysaccharides with  $\beta$ 1-4 and  $\beta$ 1-3 -  $\beta$ 1-4 linkages. The variants proteins hydrolysed CMC, lichenian, glucomannan and had highest level of enzymatic activity when assayed using lichenian (7.2 U/mg). The cellulases had very low activity on Avicel and were not active on xylan and
- 10 galactomannan. The variant cellulases had specific activities on CMC that were about 3 to about 4 times higher than the specific activity of the full-length protein (Table 1).

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181

Table 1

	Optimal temperature for catalysis (°C)	T <sub>m</sub> /T <sub>m</sub> * (°C)	specific activity on CMC (U/mg)	T <sub>1/2</sub> at 90 °C (hours)
Cel12A	>105	aggregates/120.0	0.7	3
Cel12AH	90	94.5/106.6	2.7	2
Cel12A(sp-)-H	>105	102.9/NE	3.1	5
Cel12A(sp-)	>105	103.2/120.0	3.1	Retained 80% activity after 16 hours 45 minutes at 100°C

5

NE - not examined

T<sub>m</sub> - unfolding temperature in 20 mM Tris-HCL pH7.5T<sub>m</sub>\* - unfolding temperature in 20 mM Tris-HCL pH7.5/0.5% SDS

Table 2

Sample	Relative Catalytic Activity	
	no SDS	with 0.5% SDS
Cel12A	0.7	1
$\delta$ Cel12AH	1	0.86
Cel12A(sp-)H	1	0.94

### DCS studies

Intensive aggregation of the Cel12A (expressed as fluctuations of the Cp value), starting at temperatures of 65°C - 70°C were observed in 20 mM Tris-HCl pH 7.5, and no thermal transition peaks could be detected. The variant  $\delta$ Cel12A unfolded at 94.5°C when studies under the same conditions (Fig. 5), and no aggregation was noted up to 105°C. The thermal denaturation of  $\delta$ Cel12A was irreversible. Thermal transition of CelA(sp-)H occurred at 102.9°C (Fig. 5), and thermal denaturation was irreversible. In contrast to xylanase,  $\text{Ca}^{2+}$  had no stabilizing effect on the cellulases. However, the stability of the proteins was slightly enhanced by substrate (3°C for  $\delta$ Cel12AH).

All forms of the cellulase were stabilized by 0.5 % (w/v) SDS. In addition, the detergent stabilized and prevented aggregation of full-length Cel12A (Cel12A underwent thermal transition at 120°C in the presence of 0.5% SDS). Cel12A(sp-)H and  $\delta$ Cel12AH unfolded at 120°C and 106.5°C, respectively, in presence of SDS. The thermal transitions for all tested cellulases were fully reversible when SDS was present.

Tween 20 (polyoxyethylenesorbitan monolaurate) non-ionic detergent did not stabilise the tested proteins at a concentration of 0.5 % .

### Discussion

The endo-glucanase Cel12A from *R. marinus* has been cloned and produced in  
5 *E. coli* (Halldorsdottir, S. *et al.*, *Appl Microbiol Biotechnol*, 49:277-284 (1998)).

Although the amount of protein produced in *E. coli* exceeded what could be obtained from *R. marinus* cultivations, the yield was comprehensively lower than expected. Particularly because expression was driven using a T7/lac promoter based system.

Analysis of the primary structure of Cel12A revealed an explanation for the low  
10 production yield, and interesting organisational features of the enzyme. Notably, the N-terminal contains a hydrophobic region that has features which are characteristic of signal peptides. The results of the study described herein demonstrate that this putative signal peptide is responsible for the toxicity of full-length enzyme. The optical density drop that was observed in cultures of *E. coli* when trying to express the first 38 amino  
15 acid residues of the protein, was persistent several hours after induction suggesting that no post-induction cell division occurred.

Recombinant full-length Cel12A was not stable in solution or when heated to 65° C. In addition, the optimum growth temperature of *R. marinus* (65°C) caused extensive aggregation at moderately high protein concentrations. No clear thermal  
20 transition of the full-length enzyme were observed on the calorimetric trace when the run was carried out without SDS, most likely because severe aggregation occurred before the protein was unfolded. Furthermore, a clearly visible aggregate was indeed observed when the sample was taken out of the DSC cell. However, a clear thermal transition (at 120°C) with no aggregation was observed when full-length Cel12A was  
25 tested in the presence of 0.5 % SDS. Interestingly, the specific activity of the full-length enzyme was higher in the presence of SDS. In contrast, the presence of SDS lead to a



decrease in the specific activity of the variant lacking the first 17 amino acid residues ( $\delta$ Cel12A) under the same conditions.

Cell associated glycosyl hydrolase activity has been reported earlier from *R. marinus* cultivations, where considerable activity was detected in the cell fraction (Dahlberg, L. *et al.*, *Appl. Microbiol. Biotechnol.*, 40:63-68 (1993)). To determine if Cel12A is cell associated, *R. marinus* was cultivated in the presence of CMC and the endoglucanase activity was assayed. Activity was detected in the sonicated cell fraction and in the supernatant fraction, indicating that at least some Cel12A is membrane associated.

The variants of Cel12A described herein were readily produced in large quantities by expression in *E. coli* and had improved stability and activity relative to full length Cel12A. In addition the variant proteins (e.g.,  $\delta$ Cel12AH, Cel12A(sp-)H) did not deviate substantially from the full-length enzyme in their pH optima or substrate specificities. This is consistent with the fact that the integrity of the catalytic core was maintained, and the deletion of amino-terminal amino acids did not result in appreciable changes of the theoretically estimated isoelectric point of the enzyme. However, the deletion of the first 17 amino acid residues had a pronounced effect on the stability of the enzyme in aqueous media. As discussed earlier, the majority of the full-length enzyme after the first purification step was catalytically inactive, and heating the samples at 65°C resulted in comprehensive loss of the enzyme due to aggregation. The hydrophobic residues of the putative signal peptide are likely to be solvent exposed and consequently would comprise the thermodynamic driving force for aggregation. Accordingly, heating the full-length protein can accelerate the aggregation process, since hydrophobic interactions increase with temperature in this region.

Glycosyl hydrolases are typically modular proteins, comprised of catalytic and auxiliary modules separated by recognisable linker sequences. Linker sequences occur in a variety of lengths and amino acid compositions depending on their origin (Tomme, P. *et al.*, *Cellulose Hydrolysis by Bacteria and Fungi*. in: R. K. Poole, ed., *Advances in*

*Microbial Physiology*, vol 37, London, Academic Press (1995)). Typically, linkers are rich in prolines, hydroxy amino acids, alanine or glycine, and the arrangement of these amino acids imparts flexibility necessary for their function. The putative signal peptide and the catalytic module of Cel12A are separated by a typical linker sequence. The four  
5 times repeat of glutamic acid and proline which preceding the catalytic core of the enzyme is reminiscent of the linkers that separate the different modules of the xylanase, Xyn10A from *R. marinus* (Nordberg, K.E. *et al.*, *Biochimica et Biophysica Acta*, 1353:118-124 (1997)). In addition this structure (i.e., a linker sequence that links the  
10 putative signal peptide and the catalytic module) is shared with another family 12 glycosyl-hydrolase from the hyperthermophilic archaeon *Pyrococcus furiosus* (Bauer, M.W. *et al.*, *J Bacteriology*, 181:284-290 (1999)).

The study described herein demonstrates that variant cellulases, that have the amino acid sequence of a thermostable cellulase wherein one or more amino acids which are not part of the catalytic domain of the enzyme are deleted, can be more easily  
15 produced and have improved stability and/or catalytic activity relative to certain full-length cellulases.

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the  
20 scope of the invention encompassed by the appended claims.